arXiv:1207.4053v1 [hep-ph] 17 Jul 2012

Semileptonic Transition of $\Sigma_b \to \Sigma \mu^+\mu^-$ in Family Non-universal Z' Model

N. Katırcı [*], K. Azizi [†]

Department of Physics, Doğuş University, Acıbadem-Kadıköy, 34722 Istanbul, Turkey

Abstract

Using newly available form factors obtained from light cone QCD sum rules in full theory, we study the flavor changing neutral current transition of $\Sigma_b \to \Sigma \mu^+\mu^-$ decay in the family non-universal Z' model. In particular, we evaluate the differential branching ratio, forward-backward asymmetry as well as some related asymmetry parameters and polarizations. We compare the obtained results with the predictions of the standard model. It is demonstrated that most of the mentioned observables are sensitive to family non-universal Z' gauge boson. The order of differential branching ratio shows that this decay mode can be checked at LHC in near future.

PACS numbers: 12.60.i, 12.60.Cn, 13.30.a, 13.30.Ce, 14.20.Mr

---

[*] e-mail: nkatirci@dogus.edu.tr

[†] e-mail: kazizi@dogus.edu.tr

1 Introduction

The heavy baryons containing a single heavy quark constitute a perfect laboratory to test the non-perturbative aspects of QCD. The $\Lambda_b \to \Lambda\mu^+\mu^-$ transition which is based on the flavor changing neutral current (FCNC) transition of $b \to s\mu^+\mu^-$ at quark level has been recently observed in CDF collaboration at FermiLab [1]. It is also planned to be checked at LHCb collaboration at CERN [2].

The theoretical studies on the branching ratio of $\Sigma_b \to \Sigma\ell^+\ell^-$ in standard model (SM) [3] show that this decay mode is also possible to be observed at LHC. It is expected that the experimental studies on the heavy baryons and their decay properties constitute one of the main direction of research program at LHC. Hence, the theoretical calculations can play an essential role in this regard.

Although the SM has predictions in perfect agreement with collider data up to now, there are some problems such as neutrino oscillations, baryon asymmetry, unification, dark matter, strong CP violation and the hierarchy problem, etc. which can not be addressed by the SM and still remain unsolved. To cure these deficiencies, there are a plenty of new physics (NP) models such as different extra dimension models (ED), various supersymmetric (SUSY) scenarios, etc. One of the most important new physics scenarios is Z' model, appears in many grand unified theories, such as $SU(5)$ or string-inspired $E6$ models [4–8]. The two Z' models in agenda are family non-universal Z' [9, 10] and leptophobic Z' scenarios [11, 12].

The idea of extra heavy Z boson comes from the extension of gauge group $SU(5)$, predicted by the grand unification theories to larger group $SO(10)$. The $SO(10)$ gauge group is the next important one after $SU(5)$ having one extra rank. Hence, this gauge group requires at least one extra neutral gauge boson [13]. In general, Z' gauge couplings are family universal [10, 14–19], however, due to different constructions of the different families, in string models it is possible to have family non-universal Z' couplings. In some of them, three generation of leptons and also the first and second generation of quarks have different coupling to Z' boson, when compared to the third families of quarks [10, 20, 21]. For more information about this model see for instance [9, 10, 22–26]. The leptophobic Z' model means that the new neutral gauge boson does not couple to the ordinary SM charged leptons.

The study of the Z' phenomenology is an important part of the scientific program of every present and future colliders and due to its heaviness, Z' may be used to calibrate future detectors [13]. For constraints on the mass of Z' boson and mixing parameters of this model see for example [27–29]. There are direct search for $Z' \to e^+e^-$ decay [30] at Tevatron, and the possibility to discover Z' gauge boson is analyzed in [31].

In the present work, we investigate the FCNC transition of $\Sigma_b \to \Sigma\mu^+\mu^-$ in family non-universal Z' model. In particular, we analyze the differential branching ratio, forward-backward asymmetry as well as some related asymmetry parameters and polarizations and compare with the predictions of the SM. Note that the rare baryonic $\Lambda_b \to \Lambda\ell^+\ell^-$ decay within family non-universal Z' model was analyzed in [32–34] within also family non-universal Z' model. The implications of non-universal Z' model on B meson decays were investigated in [35–43]. The $B \to K_2(\to K\pi)l^+l^-$ [44], $B \to K_1l^+l$, and $B \to K_0\pi$ [45, 46]

were investigated in the same framework as well. The effects of a family non-universal Z' gauge boson is also searched for $B \to \pi\pi$ decays in [47].

The outline of the paper is as follows. In section 2, we present the effective Hamiltonian and transition matrix elements responsible for the $\Sigma_b \to \Sigma\mu^+\mu^-$ transition. In section 3, we analyze the differential branching ratio, forward backward asymmetry, double lepton polarizations as well as some other related asymmetries in the Z' model and compare the obtained results with the SM predictions. The last section encompasses our concluding remarks.

2 The $\Sigma_b \to \Sigma\mu^+\mu^-$ transition in Family Non-universal Z' Model

2.1 The Effective Hamiltonian

Neglecting terms proportional to $\frac{V_{ub}V_{us}^*}{V_{tb}V_{ts}^*} \approx O(10^{-2})$, the effective Hamiltonian of the FCNC transition of $\Sigma_b \to \Sigma\ell^+\ell^-$, proceed via quark level $b \to s\ell^+\ell^-$ in the SM, can be written as [48–51]

$$
\begin{aligned}
\mathcal{H}^{eff} &= \frac{G_F\alpha_{em}V_{tb}V_{ts}^*}{2\sqrt{2}\pi}\Bigg[C_9^{eff}\bar{s}\gamma_\mu(1-\gamma_5)b\,\bar{\ell}\gamma^\mu\ell + C_{10}\bar{s}\gamma_\mu(1-\gamma_5)b\,\bar{\ell}\gamma^\mu\gamma_5\ell \\
&- 2m_bC_7^{eff}\frac{1}{q^2}\bar{s}i\sigma_{\mu\nu}q^\nu(1+\gamma_5)b\,\bar{\ell}\gamma^\mu\ell\Bigg] , \qquad (2.1)
\end{aligned}
$$

where α_{em} is the fine structure constant at Z mass scale, G_F is the Fermi coupling constant, V_{ij} are elements of the Cabibbo-Kobayashi-Maskawa (CKM) matrix, and C_7^{eff}, C_9^{eff}, C_{10} are the Wilson coefficients. When Z' boson is considered and $Z-Z'$ mixing is neglected, the extra part which should be added to the above effective Hamiltonian is written as [35, 36]

$$
H_{eff}^{Z'} = \frac{2G_F}{\sqrt{2}}V_{tb}V_{ts}^*\left[\frac{B_{sb}^LB_{\ell\ell}^L}{V_{tb}V_{ts}^*}\bar{s}\gamma_\mu(1-\gamma_5)b\,\bar{\ell}\gamma_\mu(1-\gamma_5)\ell + \frac{B_{sb}^LB_{\ell\ell}^R}{V_{tb}V_{ts}^*}\bar{s}\gamma_\mu(1-\gamma_5)b\,\bar{\ell}\gamma_\mu(1+\gamma_5)\ell\right] \qquad (2.2)
$$

where $B_{sb}^L = |B_{sb}^L|e^{i\varphi_s^L}$ and $B_{\ell\ell}^{L,R}$ correspond to the chiral Z' couplings to quarks and leptons, respectively. Hence, in Z' model, to get the effective Hamiltonian for the transition under consideration, we need to make the following replacements in Eq.(1) to consider Z' boson contributions besides the Z boson:

$$
\begin{aligned}
C_9^{\text{eff}} &\to C_9^{\text{eff}\prime} = C_9^{\text{eff}} - \frac{4\pi}{\alpha_s}(28.82)\frac{B_{sb}^L}{V_{tb}V_{ts}^*}(B_{\ell\ell}^L + B_{\ell\ell}^R), \\
C_{10} &\to C_{10}' = C_{10} + \frac{4\pi}{\alpha_s}(28.82)\frac{B_{sb}^L}{V_{tb}V_{ts}^*}(B_{\ell\ell}^L - B_{\ell\ell}^R), \qquad (2.3)
\end{aligned}
$$

where α_s is the strong coupling constant. Here we should mention that the Wilson coefficient C_7^{eff} remains unchanged.

In SM, the Wilson coefficient C_7^{eff} in leading logarithm approximation is written as (see [52])

$$C_7^{eff}(\mu_b) = \eta^{\frac{16}{23}} C_7(\mu_W) + \frac{8}{3}\left(\eta^{\frac{14}{23}} - \eta^{\frac{16}{23}}\right) C_8(\mu_W) + C_2(\mu_W) \sum_{i=1}^{8} h_i \eta^{a_i} \ , \tag{2.4}$$

where

$$C_2(\mu_W) = 1 \ , \quad C_7(\mu_W) = -\frac{1}{2} D_0(x_t) \ , \quad C_8(\mu_W) = -\frac{1}{2} E_0(x_t) \ . \tag{2.5}$$

The functions, $D_0(x_t)$ and $E_0(x_t)$ are defined as

$$D_0(x_t) = -\frac{(8x_t^3 + 5x_t^2 - 7x_t)}{12(1-x_t)^3} + \frac{x_t^2(2-3x_t)}{2(1-x_t)^4} \ln x_t \ , \tag{2.6}$$

$$E_0(x_t) = -\frac{x_t(x_t^2 - 5x_t - 2)}{4(1-x_t)^3} + \frac{3x_t^2}{2(1-x_t)^4} \ln x_t \ , \tag{2.7}$$

where $x_t = \frac{m_t^2}{M_W^2}$ with m_t and M_W being the top quark and W boson masses, respectively. The coefficients a_i and h_i are given as [53, 54]

$$\begin{array}{lcccccccccl} a_i = (& \frac{14}{23}, & \frac{16}{23}, & \frac{6}{23}, & -\frac{12}{23}, & 0.4086, & -0.4230, & -0.8994, & 0.1456 &), \\ h_i = (& 2.2996, & -1.0880, & -\frac{3}{7}, & -\frac{1}{14}, & -0.6494, & -0.0380, & -0.0186, & -0.0057 &). \end{array} \tag{2.8}$$

The parameter η in Eq. (2.4) is also defined as

$$\eta = \frac{\alpha_s(\mu_W)}{\alpha_s(\mu_b)} \ , \tag{2.9}$$

with

$$\alpha_s(x) = \frac{\alpha_s(m_Z)}{1 - \beta_0 \frac{\alpha_s(m_Z)}{2\pi} \ln(\frac{m_Z}{x})}, \tag{2.10}$$

where $\alpha_s(m_Z) = 0.118$ and $\beta_0 = \frac{23}{3}$.

The Wilson coefficient $C_9^{eff}(\hat{s}')$ is written as [53, 54]:

$$\begin{aligned} C_9^{eff}(\hat{s}') \ &= \ C_9^{NDR}\eta(\hat{s}') + h(z, \hat{s}')\left(3C_1 + C_2 + 3C_3 + C_4 + 3C_5 + C_6\right) \\ &\quad -\frac{1}{2} h(1, \hat{s}')\left(4C_3 + 4C_4 + 3C_5 + C_6\right) \\ &\quad -\frac{1}{2} h(0, \hat{s}')\left(C_3 + 3C_4\right) + \frac{2}{9}\left(3C_3 + C_4 + 3C_5 + C_6\right), \end{aligned} \tag{2.11}$$

where $\hat{s}' = \frac{q^2}{m_b^2}$ with $4m_l^2 \leq q^2 \leq (m_{\Sigma_b} - m_\Sigma)^2$, and

$$C_9^{NDR} \ = \ P_0^{NDR} + \frac{Y_0(x_t)}{\sin^2\theta_W} - 4Z_0(x_t) + P_E E(x_t), \tag{2.12}$$

here, NDR stands for the naive dimensional regularization scheme. We ignore the last term in this equation due to the negligible value of P_E. The $P_0^{NDR} = 2.60 \pm 0.25$ [53, 54] and the functions $Y_0(x_t)$ and $Z_0(x_t)$ are defined in the following form:

$$Y_0(x_t) = \frac{x_t}{8}\left[\frac{x_t-4}{x_t-1} + \frac{3x_t}{(x_t-1)^2}\ln x_t\right], \tag{2.13}$$

and

$$Z_0(x_t) = \frac{18x_t^4 - 163x_t^3 + 259x_t^2 - 108x_t}{144(x_t-1)^3} + \left[\frac{32x_t^4 - 38x_t^3 - 15x_t^2 + 18x_t}{72(x_t-1)^4} - \frac{1}{9}\right]\ln x_t. \tag{2.14}$$

In Eq.(2.11), the $\eta(\hat{s}')$ is given as

$$\eta(\hat{s}') \;=\; 1 + \frac{\alpha_s(\mu_b)}{\pi}\,\omega(\hat{s}'), \tag{2.15}$$

with

$$\begin{aligned}\omega(\hat{s}') \;=\; & -\frac{2}{9}\pi^2 - \frac{4}{3}\mathrm{Li}_2(\hat{s}') - \frac{2}{3}\ln\hat{s}'\ln(1-\hat{s}') - \frac{5+4\hat{s}'}{3(1+2\hat{s}')}\ln(1-\hat{s}') - \\ & \frac{2\hat{s}'(1+\hat{s}')(1-2\hat{s}')}{3(1-\hat{s}')^2(1+2\hat{s}')}\ln\hat{s}' + \frac{5+9\hat{s}'-6\hat{s}'^2}{6(1-\hat{s}')(1+2\hat{s}')}.\end{aligned} \tag{2.16}$$

At μ_b scale, for the coefficients C_j $(j = 1, ...6)$ we have

$$C_j = \sum_{i=1}^{8} k_{ji}\eta^{a_i}, \tag{2.17}$$

where k_{ji} are given as:

$$\begin{array}{llllllllll}
k_{1i} = (& 0, & 0, & \frac{1}{2}, & -\frac{1}{2}, & 0, & 0, & 0, & 0 &), \\
k_{2i} = (& 0, & 0, & \frac{1}{2}, & \frac{1}{2}, & 0, & 0, & 0, & 0 &), \\
k_{3i} = (& 0, & 0, & -\frac{1}{14}, & \frac{1}{6}, & 0.0510, & -0.1403, & -0.0113, & 0.0054 &), \\
k_{4i} = (& 0, & 0, & -\frac{1}{14}, & -\frac{1}{6}, & 0.0984, & 0.1214, & 0.0156, & 0.0026 &), \\
k_{5i} = (& 0, & 0, & 0, & 0, & -0.0397, & 0.0117, & -0.0025, & 0.0304 &), \\
k_{6i} = (& 0, & 0, & 0, & 0, & 0.0335, & 0.0239, & -0.0462, & -0.0112 &).
\end{array} \tag{2.18}$$

The other functions in Eq. (2.11) are also given as:

$$\begin{aligned}h(y,\hat{s}') \;=\; & -\frac{8}{9}\ln\frac{m_b}{\mu_b} - \frac{8}{9}\ln y + \frac{8}{27} + \frac{4}{9}x \\ & -\frac{2}{9}(2+x)|1-x|^{1/2}\begin{cases}\left(\ln\left|\frac{\sqrt{1-x}+1}{\sqrt{1-x}-1}\right| - i\pi\right), & \text{for } x \equiv \frac{4z^2}{\hat{s}'} < 1 \\ 2\arctan\frac{1}{\sqrt{x-1}}, & \text{for } x \equiv \frac{4z^2}{\hat{s}'} > 1,\end{cases}\end{aligned} \tag{2.19}$$

where $y = 1$ or $y = z = \frac{m_c}{m_b}$ and,

$$h(0, \hat{s}') \;=\; \frac{8}{27} - \frac{8}{9}\ln\frac{m_b}{\mu_b} - \frac{4}{9}\ln\hat{s}' + \frac{4}{9}i\pi. \tag{2.20}$$

We also consider the long distance contribution (Y_{LD}) coming from J/ψ family resonances to C_9^{eff} and parameterized using Breit-Wigner ansatz [48] as

$$Y_{LD} = \frac{3\pi}{\alpha_{em}^2} C^{(0)} \sum_{i=1}^{6} \kappa_i \frac{\Gamma(V_i \to l^+ l^-) m_{V_i}}{m_{V_i}^2 - q^2 - i m_{V_i}\Gamma_{V_i}}, \tag{2.21}$$

where $C^{(0)} = 0.362$, and we consider only two lowest resonances $J/\psi(1S)$ and $\psi(2S)$ and choose the corresponding phenomenological factors $\kappa_1 = 1$ and $\kappa_2 = 2$. We use the experimental results on the masses and total decay rates of dilepton decays of the considered vector charmonium states [55]. For more details about the calculation of long distance contributions, see for instance [56, 57].

Finally, the explicit expression for C_{10} is given as:

$$C_{10} = -\frac{Y_0(x_t)}{\sin^2\theta_W}\ , \tag{2.22}$$

where, $\sin^2\theta_W = 0.23$.

2.2 Transition Matrix Elements and Form Factors

The $\Sigma_b \to \Sigma\mu^+\mu^-$ decay amplitude is obtained by sandwiching the effective Hamiltonian between the initial and final states

$$\mathcal{M} = \langle \Sigma(p) | H_{\rm eff} | \Sigma_b(p+q) \rangle, \tag{2.23}$$

which reads

$$\begin{aligned}\mathcal{M} = \frac{G_F\ \alpha_{em} V_{tb}\ V_{ts}^*}{2\sqrt{2}\ \pi}\Big\{ & C_9^{\rm eff'}\ \langle \Sigma(p)|\bar{s}\gamma_\mu(1-\gamma_5)b|\Sigma_b(p+q)\rangle \bar{l}\gamma^\mu l \\ & + C_{10}'\ \langle \Sigma(p)|\bar{s}\gamma_\mu(1-\gamma_5)b|\Sigma_b(p+q)\rangle \bar{l}\gamma^\mu\gamma_5 l \\ & - 2m_b\ C_7^{\rm eff}\frac{1}{q^2}\ \langle \Sigma(p)|\bar{s}i\sigma_{\mu\nu}q^\nu(1+\gamma_5)b|\Sigma_b(p+q)\rangle \bar{l}\gamma^\mu l \Big\}\ .\end{aligned} \tag{2.24}$$

To calculate the amplitude, we need to parameterize the transition matrix elements $\langle\Sigma(p)|\bar{s}\gamma_\mu(1-\gamma_5)b|\Sigma_b(p+q)\rangle$ and $\langle\Sigma(p)|\bar{s}\sigma_{\mu\nu}q^\nu(1+\gamma_5)b|\Sigma_b(p+q)\rangle$ in terms of twelve form factors f_i, g_i, f_i^T and g_i^T $(i = 1, 2, 3)$ as follows:

$$\begin{aligned}\langle \Sigma(p)\ |\ \bar{s}\gamma_\mu(1-\gamma_5)b\ |\ \Sigma_b(p+q)\rangle &= \bar{u}_\Sigma(p)\Big[\gamma_\mu f_1(q^2) + i\sigma_{\mu\nu}q^\nu f_2(q^2) + q^\mu f_3(q^2) \\ & - \gamma_\mu\gamma_5 g_1(q^2) - i\sigma_{\mu\nu}\gamma_5 q^\nu g_2(q^2) - q^\mu\gamma_5 g_3(q^2)\Big] u_{\Sigma_b}(p+q)\ , \\ \langle \Sigma(p)\ |\ \bar{s}i\sigma_{\mu\nu}q^\nu(1+\gamma_5)b\ |\ \Sigma_b(p+q)\rangle &= \bar{u}_\Sigma(p)\Big[\gamma_\mu f_1^T(q^2) + i\sigma_{\mu\nu}q^\nu f_2^T(q^2) + q^\mu f_3^T(q^2) \\ & + \gamma_\mu\gamma_5 g_1^T(q^2) + i\sigma_{\mu\nu}\gamma_5 q^\nu g_2^T(q^2) + q^\mu\gamma_5 g_3^T(q^2)\Big] u_{\Sigma_b}(p+q)\ ,\end{aligned} \tag{2.25}$$

where u_{Σ_b} and u_Σ are spinors of Σ_b and Σ baryons, respectively.

The form factors as the main inputs in the analysis of the $\Sigma_b \to \Sigma \ell^+ \ell^-$ have been very recently calculated in full theory via light cone QCD sum rules in [3]. The fit function of transition form factors is given as

$$f_i^{(T)}(q^2)[g_i^{(T)}(q^2)] = \frac{a}{(1-\frac{q^2}{m_{fit}^2})} + \frac{b}{(1-\frac{q^2}{m_{fit}^2})^2}, \tag{2.26}$$

where the fit parameters a, b, and m_{fit} are presented in Table 1.

	a	b	m_{fit}	$q^2=0$
f_1	-0.035	0.13	5.1	0.095 ± 0.017
f_2	0.026	-0.081	5.2	-0.055 ± 0.012
f_3	0.013	-0.065	5.3	-0.052 ± 0.016
g_1	-0.031	0.15	5.3	0.12 ± 0.03
g_2	0.015	-0.040	5.3	-0.025 ± 0.008
g_3	0.012	-0.047	5.4	-0.035 ± 0.009
f_1^T	1.0	-1.0	5.4	0.0 ± 0.0
f_2^T	-0.29	0.42	5.4	0.13 ± 0.04
f_3^T	-0.24	0.41	5.4	0.17 ± 0.05
g_1^T	0.45	-0.46	5.4	-0.010 ± 0.003
g_2^T	0.031	0.055	5.4	0.086 ± 0.024
g_3^T	-0.011	-0.18	5.4	-0.19 ± 0.06

Table 1: Parameters appearing in the fit function of the form factors, f_1, f_2, f_3, g_1, g_2, g_3, f_1^T, f_2^T, f_3^T, g_1^T, g_2^T and g_3^T in full theory for $\Sigma_b \to \Sigma \ell^+ \ell^-$ together with the values of the form factors at $q^2 = 0$ [3].

3 Observables Related to the $\Sigma_b \to \Sigma \ell^+ \ell^-$ Transition

3.1 Branching Ratio

Using the amplitude mentioned above, the differential decay rate is found as (see also [58])

$$\frac{d\Gamma(z,\hat{s})}{d\hat{s}\ \ dz} = \frac{G_F^2 \alpha_{em}^2 m_{\Sigma_b}}{16384\pi^5} |V_{tb}V_{ts}^*|^2 v\sqrt{\lambda} \Bigg[\mathcal{T}_0(\hat{s}) + \mathcal{T}_1(\hat{s}) z + \mathcal{T}_2(\hat{s}) z^2 \Bigg] , \tag{3.27}$$

where $\hat{s} = \frac{q^2}{m_{\Sigma_b}^2}$ and $z = \cos\theta$ with θ being the angle between the momenta of Σ_b and ℓ^+ in the center of mass of leptons, $\lambda = \lambda(1, r, \hat{s}) = 1 + r^2 + \hat{s}^2 - 2r - 2\hat{s} - 2r\hat{s}$, $r = m_\Sigma^2 / m_{\Sigma_b}^2$ and $v = \sqrt{1 - \frac{4m_\ell^2}{q^2}}$. The functions, $\mathcal{T}_0(\hat{s})$, $\mathcal{T}_1(\hat{s})$ and $\mathcal{T}_2(\hat{s})$ are given as

$$\begin{aligned}
\mathcal{T}_0(\hat{s}) &= 32 m_\ell^2 m_{\Sigma_b}^4 \hat{s}(1 + r - \hat{s}) \Big(|D_3|^2 + |E_3|^2 \Big) \\
&+ 64 m_\ell^2 m_{\Sigma_b}^3 (1 - r - \hat{s}) \,\mathrm{Re}[D_1^* E_3 + D_3 E_1^*] \\
&+ 64 m_{\Sigma_b}^2 \sqrt{r} (6 m_\ell^2 - m_{\Sigma_b}^2 \hat{s}) \mathrm{Re}[D_1^* E_1] \\
&+ 64 m_\ell^2 m_{\Sigma_b}^3 \sqrt{r} \Big(2 m_{\Sigma_b} \hat{s} \mathrm{Re}[D_3^* E_3] + (1 - r + \hat{s}) \mathrm{Re}[D_1^* D_3 + E_1^* E_3] \Big) \\
&+ 32 m_{\Sigma_b}^2 (2 m_\ell^2 + m_{\Sigma_b}^2 \hat{s}) \Big\{ (1 - r + \hat{s}) m_{\Sigma_b} \sqrt{r} \,\mathrm{Re}[A_1^* A_2 + B_1^* B_2] \\
&- m_{\Sigma_b} (1 - r - \hat{s}) \,\mathrm{Re}[A_1^* B_2 + A_2^* B_1] - 2\sqrt{r} \Big(\mathrm{Re}[A_1^* B_1] + m_{\Sigma_b}^2 \hat{s} \,\mathrm{Re}[A_2^* B_2] \Big) \Big\} \\
&+ 8 m_{\Sigma_b}^2 \Big\{ 4 m_\ell^2 (1 + r - \hat{s}) + m_{\Sigma_b}^2 \Big[(1 - r)^2 - \hat{s}^2 \Big] \Big\} \Big(|A_1|^2 + |B_1|^2 \Big) \\
&+ 8 m_{\Sigma_b}^4 \Big\{ 4 m_\ell^2 \Big[\Sigma + (1 + r - \hat{s}) \hat{s} \Big] + m_{\Sigma_b}^2 \hat{s} \Big[(1 - r)^2 - \hat{s}^2 \Big] \Big\} \Big(|A_2|^2 + |B_2|^2 \Big) \\
&- 8 m_{\Sigma_b}^2 \Big\{ 4 m_\ell^2 (1 + r - \hat{s}) - m_{\Sigma_b}^2 \Big[(1 - r)^2 - \hat{s}^2 \Big] \Big\} \Big(|D_1|^2 + |E_1|^2 \Big) \\
&+ 8 m_{\Sigma_b}^5 \hat{s} v^2 \Big\{ - 8 m_{\Sigma_b} \hat{s} \sqrt{r} \,\mathrm{Re}[D_2^* E_2] + 4(1 - r + \hat{s}) \sqrt{r} \,\mathrm{Re}[D_1^* D_2 + E_1^* E_2] \\
&- 4(1 - r - \hat{s}) \,\mathrm{Re}[D_1^* E_2 + D_2^* E_1] + m_{\Sigma_b} \Big[(1 - r)^2 - \hat{s}^2 \Big] \Big(|D_2|^2 + |E_2|^2 \Big) \Big\},
\end{aligned} \tag{3.28}$$

$$\begin{aligned}
\mathcal{T}_1(\hat{s}) &= -16 m_{\Lambda_b}^4 \hat{s} v \sqrt{\Sigma} \Big\{ 2Re(A_1^* D_1) - 2Re(B_1^* E_1) \\
&+ 2 m_{\Lambda_b} Re(B_1^* D_2 - B_2^* D_1 + A_2^* E_1 - A_1^* E_2) \Big\} \\
&+ 32 m_{\Lambda_b}^5 \hat{s}\ v \sqrt{\Sigma} \Big\{ m_{\Lambda_b} (1 - r) Re(A_2^* D_2 - B_2^* E_2) \\
&+ \sqrt{r} Re(A_2^* D_1 + A_1^* D_2 - B_2^* E_1 - B_1^* E_2) \Big\} ,
\end{aligned} \tag{3.29}$$

$$\begin{aligned}
\mathcal{T}_2(\hat{s}) &= -8 m_{\Sigma_b}^4 v^2 \Sigma \Big(|A_1|^2 + |B_1|^2 + |D_1|^2 + |E_1|^2 \Big) \\
&+ 8 m_{\Sigma_b}^6 \hat{s} v^2 \Sigma \Big(|A_2|^2 + |B_2|^2 + |D_2|^2 + |E_2|^2 \Big) ,
\end{aligned} \tag{3.30}$$

where

$$A_1 = -\frac{2m_b}{q^2} C_7^{eff} \left(f_1^T + g_1^T \right) + C_9^{eff} \left(f_1 - g_1 \right) \ ,$$

$$
\begin{aligned}
A_2 &= A_1\,(1 \to 2)\ , \\
A_3 &= A_1\,(1 \to 3)\ , \\
B_i &= A_i\left(g_i \to -g_i;\ g_i^T \to -g_i^T\right)\ , \\
D_1 &= C_{10}\,(f_1 - g_1)\ , \\
D_2 &= D_1\,(1 \to 2)\ , \\
D_3 &= D_1\,(1 \to 3)\ , \\
E_i &= D_i\,(g_i \to -g_i)\ .
\end{aligned}
\tag{3.31}
$$

To numerically analyze the differential branching ratio with respect to q^2, we perform integral over z in the interval $z \in [-1, 1]$ in Eq. (3.27) and take the values of input parameters as $m_t = 167\ GeV$, $m_W = 80.4\ GeV$, $m_Z = 91\ GeV$, $m_b = 4.8\ GeV$, $m_c = 1.46\ GeV$, $\mu_b = 5\ GeV$, $\mu_W = 80.4\ GeV$, $|V_{tb}V_{ts}^*| = 0.041$, $G_F = 1.17 \times 10^{-5}\ GeV^{-2}$, $\alpha_{em} = \frac{1}{137}$, $m_\Sigma = 1.192\ GeV$, $m_{\Sigma_b} = 5.807\ GeV$ and $m_\mu = 0.1056\ GeV$ [55]. The other input parameters related to the family non-universal Z' model are given in Table 2, where S1 and S2 are two sets of parameters correspond to UT-fit Collaboration results [59]. These two different sets are obtained fitting the results to $\bar{B}_s - B_s$ mixing data [59]. The results of the works [45, 46] indicate that $S2$ is more favored than $S1$.

In order to maximize the effect of Z' boson, we consider the maximum values of the related parameters, i.e. $|B_{sb}^L| = 1.31 \times 10^{-3}$, $\varphi_s^{L[0]} = -79^0$, $B_{ll}^L + B_{ll}^R = -6.7 \times 10^{-2}$, and $B_{ll}^L - B_{ll}^R = -9.3 \times 10^{-2}$ for $S1$ set and $|B_{sb}^L| = 2.35 \times 10^{-3}$, $\varphi_s^{L[0]} = -86^0$, $B_{\mu\mu}^L + B_{\mu\mu}^R = -2.60 \times 10^{-2}$, and $B_{\mu\mu}^L - B_{\mu\mu}^R = -3.4 \times 10^{-2}$ for $S2$ case.

	$\left\|B_{sb}^L\right\|$	$\varphi_s^{L[0]}$	$B_{\mu\mu}^L + B_{\mu\mu}^R$	$B_{\mu\mu}^L - B_{\mu\mu}^R$
$S1$	$(1.09 \pm 0.22) \times 10^{-3}$	$^-72 \pm 7$	$(-2.80 \pm 3.90) \times 10^{-2}$	$(-6.70 \pm 2.60) \times 10^{-2}$
$S2$	$(2.20 \pm 0.15) \times 10^{-3}$	$^-82 \pm 4$	$(-1.20 \pm 1.40) \times 10^{-2}$	$(-2.50 \pm 0.90) \times 10^{-2}$

Table 2: The values of the input parameters for the Z' couplings in two sets.

We plot the differential branching ratio, forward backward asymmetry and other asymmetry parameters with respect to q^2 to check the sensitivity of these observables to $S1$ and $S2$ sets of parameters. For this aim, we plot the variation of the differential branching ratio in terms of q^2 in figure 1. From this figure, we obtain the following results:

- we see considerable discrepancies between the Z' model and the SM predictions for $S1$ set of parameters. In the case of $S2$, the difference is observed at higher values of q^2.
- The order of differential branching ratio indicates the possibility of detecting this decay mode at LHC.



Figure 1: The q^2 dependence of the differential branching ratio for the $\Sigma_b \to \Sigma\mu^+\mu^-$ decay for $S1$, and $S2$ sets of parameters as well as the SM.

3.2 Forward-backward Asymmetry

One of the most promising tools in detecting the NP effects is the forward-backward asymmetry of leptons which is defined as

$$\mathcal{A}_{FB} = \frac{N_f - N_b}{N_f + N_b}, \tag{3.32}$$

where N_f is the number of events that particle is moving ”forward” with respect to any chosen direction, and N_b is the number of events that particle moves to ”backward” direction. In technique language, the forward-backward asymmetry $\mathcal{A}_{FB}(\hat{s})$ is defined in terms of the differential decay rate as

$$\mathcal{A}_{FB}(\hat{s}) = \frac{\displaystyle\int_0^1 \frac{d\Gamma}{d\hat{s}dz}(z,\hat{s})\,dz - \int_{-1}^0 \frac{d\Gamma}{d\hat{s}dz}(z,\hat{s})\,dz}{\displaystyle\int_0^1 \frac{d\Gamma}{d\hat{s}dz}(z,\hat{s})\,dz + \int_{-1}^0 \frac{d\Gamma}{d\hat{s}dz}(z,\hat{s})\,dz} \ , \tag{3.33}$$

We depict the dependence of $\mathcal{A}_{FB}$ on q^2 in figure 2. From this figure, we conclude that the predictions of Z' model for $S1$ and $S2$ sets deviate considerably from the SM result especially at higher values of q^2.

3.3 Baryon polarization asymmetry parameter - α_Σ

Asymmetry parameters characterize the angular dependence of differential decay width for the cascade decay $\Sigma_b \to \Sigma(\to a+b)V^*(\to l^+l^-)$ with polarized and unpolarized heavy baryons. In this part, sensitivity of the baryon polarization asymmetry parameter to new Wilson coefficients is analyzed. The helicity amplitudes for the $\Sigma_b \to \Sigma\ell^+\ell^-$ decay are obtained by analyzing quasi two body decay $\Sigma_b \to \Sigma V^*$, followed by the leptonic decay of $V^* \to \ell^+\ell^-$ in [60]. These amplitudes are obtained using helicity amplitude formalism and polarization density matrix method, demonstrated in [61, 62]. Considering

$$\Sigma_b^{1/2^+} \to \Sigma^{1/2^+}\Big(\to a+b\Big) + V^*(\to \ell^+\ell^-) \ , \tag{3.34}$$



Figure 2: The q^2 dependence of the forward backward asymmetry for the $\Sigma_b \to \Sigma\mu^+\mu^-$ decay for $S1$, and $S2$ sets of parameters as well as the SM.

with V^* being off-shell γ or Z boson, the normalized joint angular decay distribution for the two cascade decay is written as [63–68]

$$\frac{d\Gamma(q^2)}{dq^2 d\cos\theta\, d\cos\theta_\Sigma} = \left|\frac{G_F\alpha_{em}}{8\sqrt{2}\pi}V_{tb}V_{ts}^*\right|^2 \frac{\sqrt{\lambda(m_{\Sigma_b}^2, m_\Sigma^2, q^2)}\sqrt{\lambda(m_\Sigma^2, m_a^2, m_b^2)}}{1024\pi^3 m_{\Sigma_b}^3 m_\Sigma^2} v\mathcal{B}(\Sigma_b \to a+b)\,|\mathcal{M}|^2 \ , \tag{3.35}$$

where $\mathcal{M}$ is calculated in [60]. For the leptonic part, in the rest frame of the intermediate boson, the angle of the anti-lepton with respect to its helicity axes is shown by θ. For the hadronic decay, in the rest frame of the Σ_b, θ_Σ is the angle of the a momentum with respect to its helicity axes.

The polar angle distribution of $\Sigma \to a+b$ decay is obtained by integrating Eq. (3.35) with respect to θ and it gives the differential decay rate in terms of q^2 and θ_Σ [60]

$$\frac{d\Gamma(q^2)}{dq^2\, d\cos\theta_\Sigma} \sim 1 + \alpha(q^2)\alpha_\Sigma(q^2)\cos\theta_\Sigma(q^2) \ , \tag{3.36}$$

where the baryon polarization asymmetry parameter α_Σ is given as

$$\begin{aligned}
\alpha_\Sigma(q^2) = {} & \frac{8}{3\Delta(q^2)}\Big\{4m_\ell^2\left|A_{+1/2,+1}\right|^2 + 2q^2\Big(\left|A_{+1/2,+1}\right|^2 + v^2\left|B_{+1/2,+1}\right|^2\Big) \\
& - 2m_\ell^2\left|A_{+1/2,0}\right|^2 - q^2\Big(\left|A_{+1/2,0}\right|^2 + v^2\left|B_{+1/2,0}\right|^2\Big) - 6m_\ell^2\left|B_{+1/2,t}\right|^2 \\
& - 4m_\ell^2\left|A_{-1/2,-1}\right|^2 - 2q^2\Big(\left|A_{-1/2,-1}\right|^2 + v^2\left|B_{-1/2,-1}\right|^2\Big) \\
& + 2m_\ell^2\left|A_{-1/2,0}\right|^2 + q^2\Big(\left|A_{-1/2,0}\right|^2 + v^2\left|B_{-1/2,0}\right|^2\Big) + 6m_\ell^2\left|B_{-1/2,t}\right|^2\Big\} \ ,
\end{aligned} \tag{3.37}$$

and the asymmetry parameter α is obtained as

$$\alpha(q^2) = \frac{8}{3\Delta(q^2)}\Big\{4m_\ell^2\left|A_{+1/2,+1}\right|^2 + 2q^2\Big(\left|A_{+1/2,+1}\right|^2 + v^2\left|B_{+1/2,+1}\right|^2\Big)$$

$$
\begin{aligned}
&+ 2m_\ell^2 \left|A_{+1/2,0}\right|^2 + 6m_\ell^2 \left|B_{+1/2,t}\right|^2 + q^2\Big(\left|A_{+1/2,0}\right|^2 + v^2 \left|B_{+1/2,0}\right|^2 \Big) \\
&- 4m_\ell^2 \left|A_{-1/2,-1}\right|^2 - 2q^2\Big(\left|A_{-1/2,-1}\right|^2 + v^2 \left|B_{-1/2,-1}\right|^2 \Big) \\
&- q^2(\left|A_{-1/2,0}\right|^2 + v^2 \left|B_{-1/2,0}\right|^2 \Big) - 2m_\ell^2 \left|A_{-1/2,0}\right|^2 - 6m_\ell^2 \left|B_{-1/2,t}\right|^2 \Big\} \ , \qquad (3.38)
\end{aligned}
$$

with

$$
\begin{aligned}
\Delta(q^2) = \frac{8}{3}\Big\{&4m_\ell^2 \left|A_{+1/2,+1}\right|^2 + 2q^2\Big(\left|A_{+1/2,+1}\right|^2 + v^2 \left|B_{+1/2,+1}\right|^2 \Big) \\
&+ 2m_\ell^2 \left|A_{+1/2,0}\right|^2 + 6m_\ell^2 \left|B_{+1/2,t}\right|^2 + q^2\Big(\left|A_{+1/2,0}\right|^2 + v^2 \left|B_{+1/2,0}\right|^2 \Big) \\
&+ 4m_\ell^2 \left|A_{-1/2,-1}\right|^2 + 2q^2\Big(\left|A_{-1/2,-1}\right|^2 + v^2 \left|B_{-1/2,-1}\right|^2 \Big) \\
&+ q^2(\left|A_{-1/2,0}\right|^2 + v^2 \left|B_{-1/2,0}\right|^2 \Big) + 2m_\ell^2 \left|A_{-1/2,0}\right|^2 + 6m_\ell^2 \left|B_{-1/2,t}\right|^2 \Big\} \ . \qquad (3.39)
\end{aligned}
$$

The definitions for A_{λ_i,λ_V} and B_{λ_i,λ_V} are given in [60] where λ_i and λ_V are helicities of lepton pairs and vector boson, respectively. We plot the dependence of the baryon asymmetry parameter α_Σ on q^2 in figure 3. From this figure, it is clear that



Figure 3: The q^2 dependence of the asymmetry parameter α_Σ.

- in general, the results obtained using two sets of parameters deviate significantly from the SM predictions.
- The effect of Z' boson on baryon asymmetry parameter for $\Sigma_b \to \Sigma\mu^+\mu^-$ channel is larger compared to that of $\Lambda_b \to \Lambda\mu^+\mu^-$ decay mode analyzed in [33].

3.4 Polar angle distribution parameters- α_θ and β_θ

The polar angle distribution of $V^* \to l^+l^-$ decay is obtained by integrating Eq. (3.35) with respect to θ_Σ which gives the differential decay rate in terms of q^2 and θ

$$
\frac{d\Gamma(q^2)}{dq^2\, d\cos\theta} \sim 1 + 2\alpha_\theta \cos\theta + \beta_\theta \cos^2\theta \ , \qquad (3.40)
$$

where

$$\alpha_\theta(q^2) = \frac{1}{\Delta_1(q^2)} 2vq^2 \mathrm{Re}\Big[A_{+1/2,+1}B^*_{+1/2,+1} - A_{-1/2,-1}B^*_{-1/2,-1}\Big] \,, \tag{3.41}$$

and

$$\begin{aligned}\beta_\theta(q^2) = \frac{1}{\Delta_1(q^2)}\Big\{ &- 4m_\ell^2\left|A_{+1/2,+1}\right|^2 + q^2\Big(\left|A_{+1/2,+1}\right|^2 + v^2\left|B_{+1/2,+1}\right|^2\Big) \\ &+ 4m_\ell^2\left|A_{+1/2,0}\right|^2 - q^2\Big(\left|A_{+1/2,0}\right|^2 + v^2\left|B_{+1/2,0}\right|^2\Big) \\ &- 4m_\ell^2\left|A_{-1/2,-1}\right|^2 + q^2\Big(\left|A_{-1/2,-1}\right|^2 + v^2\left|B_{-1/2,-1}\right|^2\Big) \\ &+ 4m_\ell^2\left|A_{-1/2,0}\right|^2 - q^2\Big(\left|A_{-1/2,0}\right|^2 + v^2\left|B_{-1/2,0}\right|^2\Big)\Big\} \,,\end{aligned} \tag{3.42}$$

with

$$\begin{aligned}\Delta_1(q^2) = &\ 4m_\ell^2\left|A_{+1/2,+1}\right|^2 + q^2\Big(\left|A_{+1/2,+1}\right|^2 + v^2\left|B_{+1/2,+1}\right|^2\Big) \\ &+ 4m_\ell^2\left|B_{+1/2,t}\right|^2 + q^2\Big(\left|A_{+1/2,0}\right|^2 + v^2\left|B_{+1/2,0}\right|^2\Big) \\ &+ 4m_\ell^2\left|A_{-1/2,-1}\right|^2 + q^2\Big(\left|A_{-1/2,-1}\right|^2 + v^2\left|B_{-1/2,-1}\right|^2\Big) \\ &+ 4m_\ell^2\left|B_{-1/2,t}\right|^2 + q^2\Big(\left|A_{-1/2,0}\right|^2 + v^2\left|B_{-1/2,0}\right|^2\Big) \,.\end{aligned} \tag{3.43}$$

We plot the dependence of the polar angle distribution parameters α_θ and β_θ on q^2 in figures 4 and 5, respectively. From these figures, we read that



Figure 4: The q^2 dependence of the asymmetry parameter α_θ.

- similar to the $\Lambda_b \to \Lambda\mu^+\mu^-$ channel considered in [33], the asymmetry parameter β_θ in Z' model gives roughly the same result as the SM; however in the case of α_θ there is a considerable discrepancy between the two model predictions.
- The zero point of α_θ parameter is shifted to the right for both sets of parameters compared to the SM prediction.
- The effect of Z' boson on the α_θ parameter in $\Sigma_b \to \Sigma\mu^+\mu^-$ channel is large compared to the $\Lambda_b \to \Lambda\mu^+\mu^-$ channel discussed in [33].



Figure 5: The q^2 dependence of the asymmetry parameter β_θ.

3.5 Double Lepton Polarizations

To analyze the double lepton polarizations, we need to define the following orthogonal unit vectors $s_i^{\pm\mu}$ with $i = L, T$ or N in the rest frame of double leptons (see for instance [69–71] for detailed information)

$$\begin{aligned}
s_L^{-\mu} &= \left(0, \vec{e}_L^{\,-}\right) = \left(0, \frac{\vec{p}_-}{|\vec{p}_-|}\right) \ , \\
s_N^{-\mu} &= \left(0, \vec{e}_N^{\,-}\right) = \left(0, \frac{\vec{p}_\Sigma \times \vec{p}_-}{|\vec{p}_\Sigma \times \vec{p}_-|}\right) \ , \\
s_T^{-\mu} &= \left(0, \vec{e}_T^{\,-}\right) = \left(0, \vec{e}_N^{\,-} \times \vec{e}_L^{\,-}\right) \ , \\
s_L^{+\mu} &= \left(0, \vec{e}_L^{\,+}\right) = \left(0, \frac{\vec{p}_+}{|\vec{p}_+|}\right) \ , \\
s_N^{+\mu} &= \left(0, \vec{e}_N^{\,+}\right) = \left(0, \frac{\vec{p}_\Sigma \times \vec{p}_+}{|\vec{p}_\Sigma \times \vec{p}_+|}\right) \ , \\
s_T^{+\mu} &= \left(0, \vec{e}_T^{\,+}\right) = \left(0, \vec{e}_N^{\,+} \times \vec{e}_L^{\,+}\right) \ ,
\end{aligned} \tag{3.44}$$

where $\vec{p}_\pm$ and $\vec{p}_\Sigma$ are the three-momenta of the leptons $\ell^\pm$ and Σ baryon. The symbols L, T and N denote longitudinal, transverse, and normal polarizations, respectively. Using the Lorentz boost, these unit vectors are transformed from the rest frame of the leptons into the center of mass (CM) frame of them along the longitudinal direction. As a result, we get

$$\left(s_L^{\mp\mu}\right)_{CM} = \left(\frac{|\vec{p}_\mp|}{m_\ell}\, , \frac{E_\ell\, \vec{p}_\mp}{m_\ell\, |\vec{p}_\mp|}\right) \ , \tag{3.45}$$

where $\vec{p}_+ = -\vec{p}_-$ and E_ℓ and m_ℓ are the energy and mass of leptons in the CM frame, respectively. Under the above transformation, the other two unit vectors, $s_N^{\pm\mu}$, $s_T^{\pm\mu}$ remain unchanged. The double lepton polarizations are given as [69–71]

$$P_{ij}(\hat{s}) = \frac{\Big(\dfrac{d\Gamma(\vec{s}_i^{\,-},\vec{s}_j^{\,+})}{d\hat{s}} - \dfrac{d\Gamma(-\vec{s}_i^{\,-},\vec{s}_j^{\,+})}{d\hat{s}}\Big) - \Big(\dfrac{d\Gamma(\vec{s}_i^{\,-},-\vec{s}_j^{\,+})}{d\hat{s}} - \dfrac{d\Gamma(-\vec{s}_i^{\,-},-\vec{s}_j^{\,+})}{d\hat{s}}\Big)}{\Big(\dfrac{d\Gamma(\vec{s}_i^{\,-},\vec{s}_j^{\,+})}{d\hat{s}} + \dfrac{d\Gamma(-\vec{s}_i^{\,-},\vec{s}_j^{\,+})}{d\hat{s}}\Big) + \Big(\dfrac{d\Gamma(\vec{s}_i^{\,-},-\vec{s}_j^{\,+})}{d\hat{s}} + \dfrac{d\Gamma(-\vec{s}_i^{\,-},-\vec{s}_j^{\,+})}{d\hat{s}}\Big)} . \tag{3.46}$$

The first (second) subindex of P_{ij} represents polarization of lepton (anti-lepton). Using the above definitions, some double lepton polarizations are obtained as

$$\begin{aligned} P_{LN}(\hat{s}) = -P_{NL}(\hat{s}) &= \frac{16\pi m_{\Sigma_b}^4 \hat{m}_\ell \sqrt{\lambda}}{\Delta'(\hat{s})\sqrt{\hat{s}}} \mathrm{Im}\Bigg\{(1-\hat{r})(A_1^* D_1 + B_1^* E_1) \\ &+ m_{\Sigma_b}\hat{s}(A_1^* E_3 - A_2^* E_1 + B_1^* D_3 - B_2^* D_1) \\ &+ m_{\Sigma_b}\sqrt{\hat{r}}\hat{s}(A_1^* D_3 + A_2^* D_1 + B_1^* E_3 + B_2^* E_1) - m_{\Sigma_b}^2 \hat{s}^2 (B_2^* E_3 + A_2^* D_3)\Bigg\} , \end{aligned} \tag{3.47}$$

$$\begin{aligned} P_{LT}(\hat{s}) &= \frac{16\pi m_{\Sigma_b}^4 \hat{m}_\ell \sqrt{\lambda} v}{\Delta'(\hat{s})\sqrt{\hat{s}}} \mathrm{Re}\Bigg\{(1-\hat{r})\Big(\left|D_1\right|^2 + \left|E_1\right|^2\Big) - \hat{s}\Big(A_1 D_1^* - B_1 E_1^*\Big) \\ &- m_{\Sigma_b}\hat{s}\Big[B_1 D_2^* + (A_2 + D_2 - D_3)E_1^* - A_1 E_2^* - (B_2 - E_2 + E_3)D_1^*\Big] \\ &+ m_{\Sigma_b}^2 \hat{s}(1-\hat{r})(A_2 D_2^* - B_2 E_2^*) \\ &+ m_{\Sigma_b}\sqrt{\hat{r}}\hat{s}\Big[A_1 D_2^* + (A_2 + D_2 + D_3)D_1^* - B_1 E_2^* - (B_2 - E_2 - E_3)E_1^*\Big] \\ &- m_{\Sigma_b}^2 \hat{s}^2 (D_2 D_3^* + E_2 E_3^*)\Bigg\} , \end{aligned} \tag{3.48}$$

$$P_{NT}(\hat{s}) = -P_{TN} = \frac{64 m_{\Sigma_b}^4 \lambda v}{3\Delta'(\hat{s})} \mathrm{Im}\Bigg\{(A_1 D_1^* + B_1 E_1^*) + m_{\Sigma_b}^2 \hat{s}(A_2^* D_2 + B_2^* E_2)\Bigg\} , \tag{3.49}$$

$$\begin{aligned} P_{NN}(\hat{s}) &= \frac{32 m_{\Sigma_b}^4}{3\hat{s}\Delta'(\hat{s})} \mathrm{Re}\Bigg\{24\hat{m}_\ell^2\sqrt{\hat{r}}\hat{s}(A_1 B_1^* + D_1 E_1^*) - 12 m_{\Sigma_b}\hat{m}_\ell^2\sqrt{\hat{r}}\hat{s}(1-\hat{r}+\hat{s})(A_1 A_2^* + B_1 B_2^*) \\ &+ 6 m_{\Sigma_b}\hat{m}_\ell^2\hat{s}\Big[m_{\Sigma_b}\hat{s}(1+\hat{r}-\hat{s})\Big(\left|D_3\right|^2 + \left|E_3\right|^2\Big) + 2\sqrt{\hat{r}}(1-\hat{r}+\hat{s})(D_1 D_3^* + E_1 E_3^*)\Big] \\ &+ 12 m_{\Sigma_b}\hat{m}_\ell^2\hat{s}(1-\hat{r}-\hat{s})(A_1 B_2^* + A_2 B_1^* + D_1 E_3^* + D_3 E_1^*) \\ &- [\lambda\hat{s} + 2\hat{m}_\ell^2(1+\hat{r}^2 - 2\hat{r} + \hat{r}\hat{s} + \hat{s} - 2\hat{s}^2)]\Big(\left|A_1\right|^2 + \left|B_1\right|^2 - \left|D_1\right|^2 - \left|E_1\right|^2\Big) \\ &+ 24 m_{\Sigma_b}^2\hat{m}_\ell^2\sqrt{\hat{r}}\hat{s}^2(A_2 B_2^* + D_3 E_3^*) - m_{\Sigma_b}^2\lambda\hat{s}^2 v^2\Big(\left|D_2\right|^2 + \left|E_2\right|^2\Big) \\ &+ m_{\Sigma_b}^2\hat{s}\{\lambda\hat{s} - 2\hat{m}_\ell^2[2(1+\hat{r}^2) - \hat{s}(1+\hat{s}) - \hat{r}(4+\hat{s})]\}\Big(\left|A_2\right|^2 + \left|B_2\right|^2\Big)\Bigg\} , \end{aligned} \tag{3.50}$$

$$P_{TT}(\hat{s}) = \frac{32 m_{\Sigma_b}^4}{3\hat{s}\Delta'(\hat{s})} \mathrm{Re}\Bigg\{ - 24\hat{m}_\ell^2\sqrt{\hat{r}}\hat{s}(A_1 B_1^* + D_1 E_1^*) - 12 m_{\Sigma_b}\hat{m}_\ell^2\sqrt{\hat{r}}\hat{s}(1-\hat{r}+\hat{s})(D_1 D_3^* + E_1 E_3^*)$$

$$
\begin{aligned}
&- 24 m_{\Sigma_b}^2 \hat{m}_\ell^2 \sqrt{\hat{r}} \hat{s}^2 (A_2 B_2^* + D_3 E_3^*) \\
&- 6 m_{\Sigma_b} \hat{m}_\ell^2 \hat{s} \Big[m_{\Sigma_b} \hat{s} (1 + \hat{r} - \hat{s}) \Big(|D_3|^2 + |E_3|^2 \Big) - 2\sqrt{\hat{r}} (1 - \hat{r} + \hat{s}) (A_1 A_2^* + B_1 B_2^*) \Big] \\
&- 12 m_{\Sigma_b} \hat{m}_\ell^2 \hat{s} (1 - \hat{r} - \hat{s}) (A_1 B_2^* + A_2 B_1^* + D_1 E_3^* + D_3 E_1^*) \\
&- [\lambda \hat{s} - 2\hat{m}_\ell^2 (1 + \hat{r}^2 - 2\hat{r} + \hat{r}\hat{s} + \hat{s} - 2\hat{s}^2)] \Big(|A_1|^2 + |B_1|^2 \Big) \\
&+ m_{\Sigma_b}^2 \hat{s} \{ \lambda \hat{s} + \hat{m}_\ell^2 [4(1 - \hat{r})^2 - 2\hat{s}(1 + \hat{r}) - 2\hat{s}^2] \} \Big(|A_2|^2 + |B_2|^2 \Big) \\
&+ \{ \lambda \hat{s} - 2\hat{m}_\ell^2 [5(1 - \hat{r})^2 - 7\hat{s}(1 + \hat{r}) + 2\hat{s}^2] \} \Big(|D_1|^2 + |E_1|^2 \Big) \\
&- m_{\Sigma_b}^2 \lambda \hat{s}^2 v^2 \Big(|D_2|^2 + |E_2|^2 \Big) \Bigg\} , \qquad (3.51)
\end{aligned}
$$

where

$$
\Delta'(\hat{s}) = \mathcal{T}_0(\hat{s}) + \frac{1}{3} \mathcal{T}_2(\hat{s}). \qquad (3.52)
$$

We plot the dependence of the double lepton polarizations on q^2 in figures $6 - 10$.



Figure 6: The q^2 dependence of Longitudinal-normal (LN) polarization of leptons



Figure 7: The same as figure 6, but for LT polarization

From these figures, we observe that



Figure 8: The same as figure 6, but for TN polarization



Figure 9: The same as figure 6, but for NN polarization



Figure 10: The same as figure 6, but for TT polarization

- in general, there are considerable discrepancies between the Z' model and the SM predictions on the double lepton polarization asymmetries under consideration for both sets of parameters except for the P_{LT}.
- The big discrepancies of the Z' model results from the SM predictions appear at lower

values of q^2 for both sets.

- In the case of P_{LN} and P_{TN}, the effect of Z' boson is large for $S2$ set of parameters. For the P_{NN} and P_{TT}, the $S1$ set of parameters is more effective than the $S2$ at lower values of q^2.

Determination of zero points of double lepton polarization asymmetries and comparison of the obtained results with the theoretical predictions can give valuable information about the existence of the Z' boson.

4 Conclusion

In the present study, we analyzed the $\Sigma_b \to \Sigma\mu^+\mu^-$ decay channel in both the SM and Z' model with two sets of parameters. Our results on the considered physical quantities overall depict considerable discrepancies between the Z' and SM models predictions. These discrepancies can be considered as signals for existing the extra Z' gauge boson. The order of the differential branching ratio indicates that this channel can be studied at LHC in near future. Any measurement on the physical quantities considered in the present work and comparison of the obtained data with the theoretical predictions can give useful information not only about the existence of the Z' gauge boson but also about the nature of participating baryons, Σ_b and Σ. Such comparison will also help us to understand which set of parameters is favored. Our numerical calculations lead that for some observables like forward-backward asymmetry, baryon polarization asymmetry parameter, polar angle distribution parameters; and P_{NN} and P_{TT} from double lepton polarization asymmetries, the $S1$ set is more effective. While for differential branching ratio; and P_{LN} and P_{TN} as other double lepton polarizations, the $S2$ set is favored. Similar analysis has been made for $\Lambda_b \to \Lambda\mu^+\mu^-$ decay channel in [33] which is in agenda of different experiments nowadays. The study of different FCNC channels can provide us with more data which may help us in searching for Z' gauge boson as new physics effect.

References

[1] T. Aaltonen et al. [CDF Collaboration], Phys. Rev. Lett. 107, 201802 (2011); arXiv:1107.3753[hep-ex].

[2] Our personal communications with Yasmine Sara Amhis from LHCb Collaboration.

[3] K. Azizi, M. Bayar, A. Ozpineci, Y. Sarac, and H. Sundu; arXiv:1112.5147[hep-ph].

[4] F. Gursey, M. Serdaroglu, Lett. Nuo. Cim. 21, (1978).

[5] G. Buchalla, G. Burdman, C. T. Hill, and D. Cominis, Phys. Rev. D 53, 5185 (1996); arXiv:hep-ph/9510376.

[6] E. Nardi, Phys. Rev. D 48, 1240 (1993); arXiv:hep-ph/9209223.

[7] J. Bernabeu, E. Nardi, and D. Tommasini, Nucl. Phys. B 409, 69 (1993); arXiv:hep-ph/9306251.

[8] V. Barger, M. Berger, and R. J. Phillips, Phys. Rev. D 52, 1663 (1995); arXiv:hep-ph/9503204.

[9] E. Eichten, I. Hinchliffe, K. D. Lane and C. Quigg, Rev. Mod. Phys. 56 579 (1984).

[10] P. langacker, and M. Plumacher, Phys. Rev. D 62 013006 (2000); arXiv:hep-ph/0001204.

[11] J. L. Lopez and D. V. Nanopoulos, Phys. Rev. D 55 397 (1997); arXiv:hep-ph/9605359.

[12] B. B. Sirvanli, Mod. Phys. Lett. A 23 347 (2008); arXiv:hep-ph/0701173.

[13] A. Leike; arXiv:hep-ph 9805494.

[14] T. K. Kuo and N. Nakagawa, Phys. Rev. D 31 1161 (1985); Phys. Rev. D 32 (1985) 306.

[15] K. K. Gan, Phys. Lett. B 209 95 (1988).

[16] E. Nardi, Phys. Rev. D 48 1240 (1993); arXiv:hep-ph/9211246.

[17] B. Holdom, Phys. Lett. B 339 114 (1994); arXiv:hep-ph/9407311.

[18] X. Zhang and B. L. Young, Phys. Rev. D 51 6584 (1995).

[19] B. Holdom and M. V. Ramana, Phys. Lett. B 365 309 (1996); arXiv:hep-ph/9509272.

[20] S. Chaudhuri, S.W. Chung, G. Hockney, and J. Lykken, Nucl. Phys. B 456 89 (1995).

[21] G. Cleaver, M. Cvetic, J. R. Espinosa, L. Everett, and P. Langacker, Nucl. Phys. B 525 3 (1998); arXiv:hep-th/9711178.

[22] Y. Zhang, Z. Cai; arXiv:1106.0163[hep-ph].

[23] A. Kundu, Phys. Lett. B 370 135 (1996); arXiv:hep-ph/9504417.

[24] J. Erler, P. Langacker, Phys. Rev. Lett. 84 212 (2000); arXiv:hep-ph/9910315.

[25] C. Caso et al., Eur. Phys. J. C3, 1 (1998).

[26] P. Langacker, Rev. Mod. Phys. 81 1199 (2008); arXiv:0801.1345 [hep-ph]

[27] K. Cheung, C. W. Chiang, N. G. Deshpande and J. Jiang, Phys. Lett. B 652 285 (2007); arXiv:hep-ph/0604223.

[28] X. G. He and G. Valencia, Phys. Rev. D 74 013011 (2006); arXiv:hep-ph/0605202.

[29] F. Abe et al. [CDF collaboration], Phys. Rev. Lett. 79 2192 (1997).

[30] A. Abulencia et al. [CDF collaboration], Phys. Rev. Lett. 96 211801 (2006); arXiv:hep-ex/0602045.

[31] M. Carena, A. Daleo, B. A. Dobrescu, T. M. P. Tait, Phys. Rev. D 70 093009 (2004); arXiv:hep-ph/0408098.

[32] A. K. Giri, R. Mohanta, Eur. Phys. J. C 45 151 (2006); arXiv:hep-ph/0510171.

[33] T. M. Aliev, M. Savci; arXiv:1202.0398[hep-ph].

[34] T. M. Aliev, M. Savci; arXiv:1202.5444[hep-ph].

[35] C. H. Chen, and H. Hatanaka, Phys. Rev. D 73 075003 (2006); arXiv:hep-ph/0602140

[36] V. Barger et. al, Phys. Lett. B 580 186 (2004); arXiv:hep-ph/0310073 ; ibid B 598, 218 (2004); arXiv:hep-ph/0406126; V. Barger et. al, Phys. Rev. D 80 055008 (2009); arXiv:(hep-ph)0902.4507.

[37] C. W. Chiang et. al, JHEP 0608 075 (2006); arXiv:hep-ph/0606122.

[38] K. Cheung et. al, Phys. Lett. B 652 285 (2007); arXiv:hep-ph/0604223.

[39] R. Mohanta, and A. K. Giri, Phys. Rev. D 79 057902 (2009); arXiv:0812.1842 [hep-ph].

[40] J. Hua, C. S. Kim, and Y. Li, Phys. Lett. B 690 508 (2010); arXiv:1002.2532 [hep-ph]; Q. Chang, Y. H. Gao, Nucl. Phys. B 845 179 (2011); arXiv:1101.1272[hep-ph].

[41] Q. Chang, X. Q. Li, and Y. D. Yang, JHEP 0905, 056 (2009); arXiv:0903.0275 [hep-ph].

[42] Y. Li, J. Hua, and K. C. Yang, JHEP 1002 082 (2010); arXiv:0907.4408 [hep-ph]; ibid 1004 052 (2010); arXiv:1002.2758 [hep-ph].

[43] S. W. Wang, G. L. Sun, X. Q. Yang, and J. S. Huang, Eur. Phys. J. C 72 1852 (2012).

[44] R. H. Li, C. D. La, W. Wang, Phys. Rev. D 83 034034 (2011).

[45] Y. Li, J. Hua, K. C. Yang, EPJC 71, 1775 (2011); arXiv:1107.0630[hep-ph].

[46] Y. Li, X. J. Fan, J. Hua, E. L. Wang; arXiv:1111.7153[hep-ph].

[47] Q. Chang, X. Q. Li, Y. D. Yang, Int. J. Mod. Phys. A 26 1273 (2011); arXiv:1003.6051v2 [hep-ph].

[48] G. Buchalla, A. J. Buras, M. E. Lautenbacher, Rev. Mod. Phys. 68 1125 (1996); arXiv:hep-ph/9512380.

[49] G. Bobeth, A. J. Buras, F. Kruger, J. Urban, Nucl. Phys. B 630, 387 (2002); arXiv:hep-ph/0112305.

[50] W. Altmannshofer, P. Ball, A. Bharucha, A. Buras, D. Straub, M. Wick, JHEP 0901,019 (2009); arXiv:0811.1214.

[51] A. Ghinculov, T. Hurth, G. Isidori, Y. P. Yao, Nucl. Phys. B 685, 351 (2004); arXiv:hep-ph/0312128.

[52] A. Buras, M. Misiak, M. Münz and S. Pokorski, Nucl. Phys. B 424, 374 (1994); arXiv:hep-ph/9311345.

[53] M. Misiak, Nucl. Phys. B 393, 23 (1993); Erratum ibid B 439, 161 (1995).

[54] A.J. Buras, M. Münz, Phys. Rev. D 52, 186 (1995); arXiv:hep-ph/9501281.

[55] J. Beringer et al. (Particle Data Group), Phys. Rev. D 86, 010001 (2012).

[56] M. Beneke, G. Buchalla, M. Neubert and C.T. Sachrajda, Eur. Phys. J. C 61 439 (2009);arXiv:0902.4446[hep-ph].

[57] A. Khodjamirian, Th. Mannel , A. A. Pivovarov and Y. M. Wang; arXiv:1006.4945[hep-ph].

[58] T. M. Aliev, K. Azizi, M. Savci, Phys. Rev. D 81 056006 (2010); arXiv:1001.0227[hep-ph].

[59] M. Bona et. al [UT-fit Collaboration], PMC Phys. A 3 6 (2009); arXiv:0803.0659 [hep-ph].

[60] T. M. Aliev, M. Savcı, JHEP 0605 001 (2006); arXiv:hep-ph/0507324.

[61] P. Bialas, J. G. Körner, M. Krömer, and K. Zalewski, Z. Phys. C 57 115 (1993); arXiv:hep-ph/9303208.

[62] J. G. Korner and M. Kromer, Phys. Lett. B 275 495 (1992).

[63] P. Bialas, J. G. Körner, M. Krämer, and K. Zalewski, Z. Phys. C 57 (1993) 115;

[64] T. Mannel and G. A. Schuler, Phys. Rev. D 279, 194 (1992).

[65] M. Tanaka Phys. Rev. D 47 4969 (1993).

[66] M. Gremm, G. Köpp, and L. M. Sehgal, Phys. Rev. D 52 1588 (1995); arXiv:hep-ph/9502207.

[67] C. S. Huang and H. G. Yan, Phys. Rev. D 56, 5981 (1997).

[68] J. G. Körner and D. Pirjol, Phys. Lett. B 34 399 (1994); arXiv:hep-ph/9405360.

[69] T. M. Aliev, M. Savci, B. B. Sirvanli, Eur. Phys. J. C 52, 375 (2007), arXiv:hep-ph/0608143.

[70] T. M. Aliev, V. Bashiry, M. Savci, Eur. Phys. J. C 38, 283 (2004), arXiv:hep-ph/0409275.

[71] W. Bensalem, D. London, N. Sinha and R. Sinha, Phys. Rev. D 67, 034007 (2003), arXiv:hep-ph/0209228.